

Quantum Limit of Deterministic Theories

Massimo BLASONE,^{1,3,*} Petr JIZBA^{2,†} and Giuseppe VITIELLO^{3,‡}

¹*Theoretical Physics, Blackett Laboratory, Imperial College London, SW7 2AZ London, U.K.*

²*Institute of Theoretical Physics, University of Tsukuba, Ibaraki 305-8571, Japan*

³*Dipartimento di Fisica, INFN and INFN, Università di Salerno, I-84100 Salerno, Italy*

(Received February 3, 2003)

We show that the quantum linear harmonic oscillator can be obtained in the large N limit of a classical deterministic system with $SU(1, 1)$ dynamical symmetry. This is done in analogy with recent work by G. 't Hooft who investigated a deterministic system based on $SU(2)$. Among the advantages of our model based on a non-compact group is the fact that the ground state energy is uniquely fixed by the choice of the representation.

KEYWORDS: 't Hooft's quantization, large N limit, $SU(1, 1)$ symmetry

1. Introduction

Large N limit, has long been recognized, plays a fundamental rôle in obtaining classical limit of various quantum systems. Quantum spin models,¹⁾ quantum vector models²⁾ and $U(N)$ lattice gauge theories³⁾ provide examples. This observation has further been reinforced by numerical evidences⁴⁾ showing that the dynamical phase transitions in many non-integrable quantum field models in the large N limit can be identified with the stochastic transitions from regular to chaotic motion in the corresponding classical systems.

Yet, recently a reverse rôle of the large N limit has been conjectured by G. 't Hooft,^{5,6)} i.e., the large N limit of a deterministic theory may give rise to a genuine quantum system. This often happen in conjunction with information loss.^{5,7)} 't Hooft's work has aroused substantial interest in the possibility of obtaining a whole range of quantum models from purely classical considerations. It should be stressed that due to the non-local nature of the information loss and due to emergence of (non-local) geometric phases, Bell's inequalities cannot be utilized. The above scenario has been studied in numerous deterministic systems.⁶⁻¹⁰⁾

't Hooft's conjecture brings a new perspective in the understanding of the connection between classical chaotic dynamics and quantum mechanics. It has been recently shown,⁹⁾ that quantum gauge field theory can emerge in the infrared limit of a higher-dimensional, classical (non-Abelian) gauge field theory, known to have chaotic behavior.

The purpose of this paper is to provide a concrete example illustrating the basic features of 't Hooft's proposal on a simple deterministic system with a *non-compact* dynamical group. Our model is alternative to the $SU(2)$ model given by 't Hooft^{5,6)} and presents some advantages with respect to it in the fact that the non-compact $SU(1, 1)$ structure better fits the Weyl–Heisenberg algebra of the quantum linear harmonic oscillator (LHO), to which our system reduces in the large N limit.

In Sec. 2 we briefly review 't Hooft's $SU(2)$ system. In Sec. 3 we introduce the $SU(1, 1)$ deterministic system¹⁰⁾ and in Sec. 4 the corresponding quantum limit to LHO is performed. Sec. 5 is devoted to concluding remarks.

2. 't Hooft's $SU(2)$ Model and Its Quantum Limit

Let us briefly recapitulate 't Hooft's $SU(2)$ example.⁶⁾ This consists of an autonomous dynamical system represented by N distinct states (k), $k = 1, \dots, N$. The time evolution takes place in discrete time steps of equal size, $\Delta t = \tau$ with periodicity condition (k) = ($k + N$). By admitting the following representation

$$(0) = (N) = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}; \dots; (N-1) = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad (1)$$

the evolution is regulated by the Hamiltonian H as:

$$U(\tau) = e^{-iH\tau} = e^{-i\frac{\pi}{N}} \begin{pmatrix} 0 & & & 1 \\ 1 & 0 & & \\ & 1 & 0 & \\ & & \ddots & \ddots \\ & & & 1 & 0 \end{pmatrix}. \quad (2)$$

The factor $-i\pi/N$ is 't Hooft's phase choice. Evolution matrix satisfies the condition $U^N = -\mathbb{1}$ and hence its (un-normalized) eigenstates, say $|n\rangle$, $n = 0, 1, \dots, N-1$ are

$$|n\rangle = \sum_{k=1}^N e^{i2\pi nk/N} (k). \quad (3)$$

By defining $N \equiv 2l+1$ and $n \equiv m+l$ (i.e., $m = -l, \dots, l$) 't Hooft mapped his system onto $SU(2)$ algebra as follows

$$H |l, m\rangle = \omega(n + \frac{1}{2}) |l, m\rangle. \quad (4)$$

$$L_3 |l, m\rangle = m |l, m\rangle,$$

$$L_+ |l, m\rangle = \sqrt{(2l-n)(n+1)} |l, m+1\rangle,$$

* e-mail: m.blasone@imperial.ac.uk

† e-mail: petr@cm.ph.tsukuba.ac.jp

‡ e-mail: vitiello@sa.infn.it

$$L_- |l, m\rangle = \sqrt{(2l - n + 1)n} |l, m - 1\rangle. \quad (5)$$

Here $\omega = 2\pi/N\tau$. The continuous limit is obtained by letting $l \rightarrow \infty$ and $\tau \rightarrow 0$ with ω fixed.

As shown in ref. 10 such a limit corresponds to a *group contraction*, and by defining $a^\dagger \equiv L_+/\sqrt{2l}$ and $a \equiv L_-/\sqrt{2l}$ one recovers for $l \rightarrow \infty$ and ω fixed the Weyl-Heisenberg algebra $h(1)$ of quantum LHO, i.e.,

$$\begin{aligned} H |n\rangle &= \omega(n + 1/2) |n\rangle, \\ a^\dagger |n\rangle &= \sqrt{n+1} |n+1\rangle, \quad a |n\rangle = \sqrt{n} |n-1\rangle. \end{aligned} \quad (6)$$

't Hooft's system may be mimicked, for instance, by a charged particle in a cylindrical magnetron*. Then the particle trajectory is basically a cycloid wrapped around the center of the magnetron. The actual parameters (and qualitative nature) of the cycloid are specified by the Larmor frequency $\omega_L = qB/2m$. To implement the discrete time evolution we confine ourself only to the observation of the largest radius positions of the particle. So particularly we disregard any information concerning the actual underlying trajectory. This corresponds to loss of information. If the orbital frequency is an integer multiple of Larmor frequency then the particle proceeds via discrete time evolution with $\tau = 2\pi/\omega_L$ and returns into its initial position after one revolution, see Fig. 1. The continuous limit then corresponds to an appropriate increase in magnetic field.




Fig. 1. 't Hooft's deterministic system for $N = 7$. An underlying continuous dynamics is introduced, where $x(t) = \cos(\alpha t) \cos(\beta t)$ and $y(t) = -\cos(\alpha t) \sin(\beta t)$. At the times $t_j = j\pi/\alpha$, with j integer, the trajectory touches the external circle, i.e., $R^2(t_j) = x^2(t_j) + y^2(t_j) = 1$, and thus π/α is the frequency of the discrete ('t Hooft) system. At time t_j , the angle of $R(t_j)$ with the positive x axis is given by: $\theta_j = j\pi - \beta t_j = j(1 - \beta/\alpha)\pi$. When β/α is a rational number, say q , the system returns to the origin (modulo 2π) after N steps. To ensure that the N steps cover only one circle, we have to impose $\alpha(t_j) = j 2\pi/N$. Thus, in order to reproduce 't Hooft's system for $N = 7$ we choose $q = 5/7$.

3. The $SU(1, 1)$ Deterministic System

Let us now consider a different deterministic system, which can be related to the $SU(1, 1)$ algebraic structure.




Fig. 2. A deterministic system based on $SU(1, 1)$ and obtained by using the same dynamics as in Fig. 1 with $\beta/\alpha = 5/3 + \pi/40$.

The system in question consists of two sub-systems, each of them comprised of a particle moving along a circle in discrete equidistant jumps. Both particles and circle radii might be different the only common constraint is that particles are synchronized in their jumps. Let us further assume that for the two particles the circumference and the elementary jump lengths are *incommensurable* so that particles never come back into the original position after a finite number of jumps. We will label the corresponding states (positions) as $(n)_A$ and $(n)_B$. The synchronized time evolution materializes via discrete and identical time steps $\Delta t = \tau$ as follows:

$$\begin{aligned} t &\rightarrow t + \tau; & (1)_A &\rightarrow (2)_A \rightarrow (3)_A \rightarrow (4)_A \dots, \\ & & (1)_B &\rightarrow (2)_B \rightarrow (3)_B \rightarrow (4)_B \dots \end{aligned}$$

This evolution is, of course, completely deterministic. In fact, it is not difficult to devise a gedanken experiment producing such an evolution. Using again the magnetron system and assuming that the Larmor frequency and orbital frequency are incommensurable then particles A and B "return" into their initial positions after infinitely revolutions, see Fig. 2. Let us regularize the motion by assuming that particles come after N revolutions back into origin: N is our limiting (cutoff) parameter. Representing the actual states (positions) $(n)_{A,B}$ as in eq. (1) then the one-time-step evolution operator acting on $(n)_A \otimes (m)_B$ reads

$$\begin{aligned} U(\tau) &\equiv e^{-iH\tau} = e^{-iH_A\tau} \otimes e^{-iH_B\tau} \\ &= \begin{pmatrix} 0 & 0 & \dots & 1 \\ 1 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots \\ & & \ddots & \ddots \end{pmatrix}_A \otimes \begin{pmatrix} 0 & 0 & \dots & 1 \\ 1 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots \\ & & \ddots & \ddots \end{pmatrix}_B. \end{aligned}$$

Note that the first lines in the matrices describe the "spurious" evolution

$$\begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}_A \otimes \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}_B \rightarrow \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}_A \otimes \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}_B. \quad (7)$$

Such an evolution ensures that $U(\tau)$ is unitary and compatible with the evolution on Fig. 2.

* Cylindrical magnetron is a device with a radial, cylindrically symmetric electric field that has in addition a perpendicular uniform magnetic field.

By diagonalizing $U(\tau)$ we obtain that the two matrices involved in $U(\tau)$ have eigenvalues $\lambda_n = e^{i2\pi n/N}$ ($n = 0, \dots, N-1$) with multiplicity 1. For future convenience it is important to realize that

$$\lambda_n = e^{i2\pi n/N} = e^{i2\pi \zeta n},$$

where we have defined $\zeta = (1 - N)/N$. So there is a basis (polar basis) in which $U(\tau)$ has a diagonal form:

$$U(\tau) = U_A(\tau) \otimes U_B(\tau), \quad (8)$$

$$U_{A,B}(\tau) \equiv \text{diag} \left(1, e^{i2\pi \zeta}, \dots, e^{i2\pi \zeta (N-1)} \right)_{A,B}.$$

Let us denote the corresponding eigenstates of $U(\tau)$ as $\{|n_A\rangle \otimes |m_B\rangle; n_A, m_B \in [0, (N-1)]\}$ so that

$$U_A(\tau)|n_A\rangle = e^{i2\pi \zeta n_A}|n_A\rangle, \quad (9)$$

$$U_B(\tau)|m_B\rangle = e^{i2\pi \zeta m_B}|m_B\rangle. \quad (10)$$

We observe that with respect to the original non-polar basis (i.e., position vectors) the (un-normalized) eigenvectors read

$$|n_A\rangle = \sum_{k=1}^N e^{-i2\pi \zeta n_A k} (k)_A, \quad (11)$$

$$|m_B\rangle = \sum_{k=1}^N e^{-i2\pi \zeta m_B k} (k)_B. \quad (12)$$

It should be stressed that there is a deep qualitative difference between $(\dots)$ and $|\dots\rangle$ vectors. The first describe the time dependent set of states characterizing the deterministic evolution. The second describe the time independent states (eigenstates of the formal Hamiltonian, see below) which have no connection with actual particle trajectory. As a result of (9) and (10) we have

$$|n_A\rangle \otimes |m_B\rangle = \sum_{k,l=1}^N e^{-i2\pi \zeta (n_A k + m_B l)} (k)_A \otimes (l)_B. \quad (13)$$

Defining $(n_A - m_B)/2 \equiv j$ (i.e., $|j| = 0, 1/2, 1, 3/2, \dots, (N-1)/2$) and $(n_A + m_B)/2 \equiv m$ (i.e., $m = |j|, |j| + 1, \dots, (N-1)$), we may pass from the basis $|n_A\rangle \otimes |m_B\rangle$ to $|j, m\rangle$ basis. It is then clear that

$$|j, m\rangle = \sum_{l,k=1}^N e^{-i2\pi \zeta (m(k+l) + j(k-l))} (k)_A \otimes (l)_B. \quad (14)$$

Correspondingly,

$$\begin{aligned} U(\tau)|j, m\rangle &= U_A(\tau) \otimes U_B(\tau)|j, m\rangle \\ &= \sum_{k,l}^N e^{-i2\pi \zeta (m(k+l) + j(k-l))} U_A(\tau)(k)_A \otimes U_B(\tau)(l)_B \\ &= \sum_{k,l=1}^N e^{i2\pi \zeta (m(k+l-2) + j(k-l))} (k)_A \otimes (l)_B \\ &= e^{i2\pi \zeta 2m} |j, m\rangle. \end{aligned} \quad (15)$$

Similarly,

$$U_A(\tau) \otimes U_B(-\tau) |j, m\rangle = e^{i2\pi \zeta 2j} |j, m\rangle. \quad (16)$$

We note that (15) can be equivalently written as

$$U(\tau)|j, m\rangle = e^{i2\pi \zeta (2m-n)} |j, m\rangle. \quad (17)$$

Here n is an arbitrary integer allowed by the identity $e^{i2\pi n} = 1$.

4. Quantum Limit and Zero-Point Energy

Equation (17) implies that the total Hamiltonian has the spectrum

$$\begin{aligned} H |j, m\rangle &= (H_A + H_B) |j, m\rangle = \frac{2\pi}{\tau} (-\zeta 2m + n) |j, m\rangle \\ &= \omega [(n_A + m_B) - n/\zeta] |j, m\rangle, \end{aligned} \quad (18)$$

which in the $N \rightarrow \infty$ limit is

$$H |j, m\rangle = \frac{2\pi}{\tau} (n_A + m_B + n) |j, m\rangle = \omega (2m + n) |j, m\rangle.$$

From dimensionality reasons we can view $\omega = -\zeta 2\pi/\tau$ as a formal frequency. Note that ω is finite in the large N limit; $\omega \rightarrow 2\pi/\tau$. Let us now define:

$$L_3 \equiv \frac{\mathcal{H}}{\omega} \equiv \frac{(H_A + H_B)}{2\omega}; \quad \mathcal{C} \equiv \frac{(H_A - H_B)}{2\omega}. \quad (19)$$

Equation (16) implies that

$$(H_A - H_B) |j, m\rangle = -\frac{2\pi \zeta}{\tau} 2j |j, m\rangle, \quad (20)$$

and for $N \rightarrow \infty$ we have

$$\frac{(H_A - H_B)}{2} |j, m\rangle = \omega j |j, m\rangle. \quad (21)$$

Note that $(H_A - H_B)$ represents the energy excess of the system A over system B and hence it is the system (dynamical) invariant (i.e., among others, it commutes with H).

Applying the $N \rightarrow \infty$ limit and setting $n = 1$ we can map our deterministic system onto $SU(1, 1)$ algebra in the following way:

$$L_3 |j, m\rangle = (m + 1/2) |j, m\rangle, \quad \mathcal{C} |j, m\rangle = j |j, m\rangle. \quad (22)$$

To close the algebra we need to construct the ladder operators L_+ and L_- in terms of the system variables. This can be done by the following observation: We define the position operators (or time pointers)

$$N_A (k)_A = k(k)_A, \quad ; \quad N_B (l)_B = l(l)_B, \quad (23)$$

then

$$e^{-i2\pi \zeta (N_A + N_B)} |j, m-1\rangle = |j, m\rangle. \quad (24)$$

This in turn suggests that

$$\begin{aligned} L_+ &= e^{-i2\pi \zeta (N_A + N_B)} \left(\sqrt{(L_3 + 1/2)^2 - \mathcal{C}^2} \right), \\ L_- &= \left(\sqrt{(L_3 + 1/2)^2 - \mathcal{C}^2} \right) e^{i2\pi \zeta (N_A + N_B)} \end{aligned}$$

$$= e^{i2\pi\zeta(N_A+N_B)} \left(\sqrt{(L_3 - 1/2)^2 - \mathcal{C}^2} \right), \quad (25)$$

or in other words positive times (positions) ($N_A > 0$, $N_B > 0$) are responsible for the forward ladder operator and negative times (positions) ($N_A < 0$, $N_B < 0$) are responsible for the backward ladder operator. As $\mathcal{C}$ clearly commutes with L_- , L_+ and L_3 it coincides with the Casimir operator of $SU(1, 1)$ algebra. Eigenvalues of the Casimir set, as usually, the algebra representation.

Some comments are in order. Since the $SU(1, 1)$ group is well known (see e.g., ref. 12), we only recall that it is locally isomorphic to the (proper) Lorentz group in two spatial dimensions $SO(2, 1)$ and it differs from $SU(2)$ only in a sign in the commutation relation: $[L_+, L_-] = -2L_3$. $SU(1, 1)$ representations are well known. In particular, if we define $n \equiv m - |j|$ and $k \equiv |j| + 1/2$ we obtain the $SU(1, 1)$ discrete series D_k^+

$$\begin{aligned} L_3 |k, n\rangle &= (n + k) |k, n\rangle, \\ L_+ |k, n\rangle &= \sqrt{(n + 2k)(n + 1)} |k, n + 1\rangle, \\ L_- |k, n\rangle &= \sqrt{(n + 2k - 1)n} |k, n - 1\rangle. \end{aligned} \quad (26)$$

Note that the basic parameters of A and B systems (such as mass of particles or radii of circles) determine the value of $(H_A - H_B)$ and hence the representation of $SU(1, 1)$. So particularly when both systems A and B are identical, then $\mathcal{H}$ describes the energy of the single system (be it A or B). The existence of the second system is then imprinted via the choice of the representation of $SU(1, 1)$. In fact, in this particular situation we have $j = 0$ which corresponds to the fundamental representation $D_{1/2}^+$.

By identifying $|\frac{1}{2}, n\rangle \equiv |n\rangle$ one has

$$\begin{aligned} L_3 |n\rangle &= (n + 1/2) |n\rangle, \\ L_+ |n\rangle &= (n + 1) |n + 1\rangle; \quad L_- |n\rangle = n |n - 1\rangle. \end{aligned} \quad (27)$$

We note that the ladder operators matrix elements do not carry the square roots, as on the contrary happens in the usual Weyl–Heisenberg algebra $\mathfrak{h}(1)$ of LHO. In order to connect the system eqs. (27) with the quantum LHO we introduce the following mapping in the universal enveloping algebra of $SU(1, 1)$:

$$a = \frac{1}{\sqrt{L_3 + 1/2}} L_-; \quad a^\dagger = L_+ \frac{1}{\sqrt{L_3 + 1/2}}, \quad (28)$$

which gives us the wanted $\mathfrak{h}(1)$ structure (6), with $\mathcal{H} = \omega L_3$. So we have found one-to-one (non-linear) mapping between the deterministic system represented in terms of the $SU(1, 1)$ and the quantum harmonic oscillator. The corresponding mapping (28) is nothing but the non-compact analog¹³⁾ of the well-known Holstein–Primakoff representation for $SU(2)$ spin systems.¹⁴⁾ In ref. 10 this non-linear mapping has been related to the canonical formalism for quantum dissipation.¹¹⁾

We finally remark that the $1/2$ term in the L_3 eigenvalues is implied by the used representation. Moreover, after a period $T = 2\pi/\omega$, the evolution of the state presents a phase π that it is not of dynamical origin: $e^{-iHT} \neq 1$, it is a geometric phase. Thus the zero point energy is

strictly related to this geometric phase (which confirms the result of refs. 7). We also observe that in order to recover the original state, one must perform one more cycle. This is related to the isomorphism between $SO(2, 1)$ and $SU(1, 1)/Z_2$ and therefore it is $e^{i2 \times 2\pi L_3} = 1$.

5. Conclusions

The main objective of this paper was to discuss the algebraic structure underlying the “quantum limit” procedure recently proposed by G. ’t Hooft.⁶⁾

We have shown that the large N limit prescription used there for obtaining truly quantum systems out of deterministic ones has exemplified in our case by removing the cutoff in the regularized algebraic description of two synchronized charged particles moving in a cylindrical magnetron.

When the cutoff was taken to infinity we could recognize that the algebraic underpinning of our deterministic system was that of $SU(1, 1)$. The fundamental representation $D_{1/2}^+$ then corresponded to the situation when the second particle was “forgotten”. In this case we were able to find a one-to-one mapping of our $SU(1, 1)$ deterministic system onto the quantum LHO. Such a mapping is an analog of the well known Holstein–Primakoff mapping for diagonalizing the ferromagnet Hamiltonian.¹⁴⁾

Acknowledgments

The authors thank the ESF network COSLAB, MIUR, INFN, INFM, EPSRC and JSPS for partial support.

References

- 1) B. Simon: Commun. Math. Phys. **71** (1980) 247; E. Lieb: Commun. Math. Phys. **31** (1973) 327; R. Shankar: Phys. Rev. Lett. **45** (1980) 1088.
- 2) F. A. Berezin: Commun. Math. Phys. **63** (1978) 131.
- 3) L. G. Yaffe: Rev. Mod. Phys. **54** (1982) 407.
- 4) T. Prosen: Phys. Rev. Lett. **80** (1998) 1808; T. Prosen: Phys. Rev. E **60** (1999) 1658; Phys. Rev. E **60** (1999) 3949.
- 5) G. ’t Hooft: in *Basics and Highlights of Fundamental Physics* (Erice, 1999) [hep-th/0003005 (2000)].
- 6) G. ’t Hooft: hep-th/0104080 (2001); hep-th/0105105 (2001); quant-ph/0212095 (2002).
- 7) M. Blasone, P. Jizba and G. Vitiello: Phys. Lett. A **287** (2001) 205; M. Blasone and P. Jizba: Can. J. Phys. **80** (2002) 645; quant-ph/0102128 (2001).
- 8) C. van de Bruck: gr-qc/0001048 (2000).
- 9) T. S. Biró, B. Müller and S. G. Matinyan: Found. Phys. Lett. **14** (2001) 471; hep-th/0301131 (2003).
- 10) M. Blasone, E. Celeghini, P. Jizba and G. Vitiello, Phys. Lett. A **310** (2003) 393.
- 11) E. Celeghini, M. Rasetti and G. Vitiello: Ann. Phys. **215** (1992) 156.
- 12) A. M. Perelomov: *Generalized Coherent States and Their Applications* (Springer, Berlin, 1986).
- 13) C. C. Gerry: J. Phys. A **16** (1983) L1.
- 14) T. Holstein and H. Primakoff: Phys. Rev. **58** (1940) 1098; M. N. Shah, H. Umezawa and G. Vitiello: Phys. Rev. B **10** (1974) 4724.